

Gulf of Mexico Harmful Algal Bloom Bulletin


Region: Texas

Monday, 11 April 2016

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service Last bulletin: Monday, April 4, 2016



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from April 1 to 8: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/hab_publication/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

Conditions Report

Karenia brevis (commonly known as Texas red tide) ranges from not present to background concentrations along the coast of Texas. No respiratory irritation is expected Monday, April 11 through Monday, April 18.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Data from Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, indicates that *Karenia brevis* concentrations ranges from 'not present' to 'background' (TAMU; 4/5-8). For information on area shellfish restrictions, contact the Texas Department of State Health Services.

Recent MODIS Aqua imagery (4/10, shown left) is obscured by clouds in patches alongand offshore the Texas coast from Freeport to south of the Rio Grande, limiting analysis. Elevated to high chlorophyll (3-15 μ g/L) is visible along- and offshore the coast of Texas from Sabine Pass to Freeport. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 10 km north from the Port Aransas region from April 10-14.

Yang, Lalime




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

Wind Analysis

Port Aransas to Matagorda Ship Channel: South winds (10-20kn, 5-10m/s) today becoming southwest (5-10kn, 3-5m/s) after midnight. Northeast to east winds (5-15kn, 3-8m/s) Tuesday through Friday night.

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive: http://tidesandcurrents.noaa.gov/hab/bulletins.html



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).